## Practice Problems for Biochemistry, Session 5: Biochemical Reactions, Enzymes and ATP


## **Question 1**

You are looking at the following biological reactions that are catalyzed by specific enzymes as shown below.

Compound A ◀	E1	Compound B
Compounds C + D◀	E2	→ Compound E
Compound F ◀	E3	——▶Compound G

Note:

- In Reaction #1, Enzyme E1 catalyzes the conversion of Compound A to Compound B. This reaction is truly reversible and the reverse reaction is as likely as the forward reaction.
- In Reaction #2, Enzyme E2 catalyzes an energy requiring reaction that uses Compounds C and D as substrates to produce Compound E.
- In Reaction #3, Enzyme E3 catalyzes an energy producing reaction that converts Compound F to Compound G.
- a) Draw the energy profiles of reactions #1, #2 and #3 on the axes given below. On the diagram, please label the reactants and the products.



- b) You decide to reproduce Reaction #2 in three test tubes. You may assume that this biological reaction occurs at a 37°C and a pH of 7.4 under normal conditions.
  - Test tube #1: You perform the reaction at 70°C and a pH of 7.4 and observe that no Compound E is produced. When the temperature is brought to 37°C, Compound E is produced at a rate similar to normal condition.
  - **Test tube #2:** You perform the reaction at **37°C** and a **pH of 10.4** and observe that no Compound E is produced. When the pH is brought to **7.4**, Compound E is produced at a rate similar to normal condition.
  - Test tube #3: You perform the reaction at 37°C and a pH of 7.4 in the presence of proteases and observe that no Compound E is produced. The effect of protease treatment persists even after its removal from reaction mixture.

Explain the effect of the changed reaction parameters in each of the above test tubes on **structure** and **function** of Enzyme E2.


Reaction parameters	Effect on Enzyme E2	Is the effect reversible/ irreversible?
70°C in test tube #1		
10.4 pH in test tube #2		
Protease in Test tube #3		

c) An enzyme may alter some or all of the following parameters. For each choose whether the enzyme would "increase" or "decrease" that parameter. If the enzyme does not affect that parameter, write "no change".

Parameter	Effect of enzyme
Free energy of the products	
Rate of forward reaction	
Rate of reverse reaction	
Energy of activation	
Overall free energy change	

## Question 2

- a) On the axes below, draw and label the products and reactants,  $\Delta G$ , and energy of activation for reaction 1 below.
  - 1) A  $\Longrightarrow$  B, where the forward reaction has a  $\Delta G = -3$ , and an Energy of activation = +0.5



Compare reaction 1 (above) to reactions 2 and 3 (below).

- 2) C  $\rightarrow$  D, where the forward reaction has a  $\Delta G$  = +2, and an Energy of activation = +2.5 F, where the forward reaction has a  $\Delta G$  = +0.3, and an Energy of activation = +2
- b) Which of these reactions, 1, 2, or 3 is most likely to proceed in the forward direction in the absence of an enzyme? If you do not have enough information to answer this question, write "Can't tell" below. Explain your answer.
- c) Which of these reactions, 1, 2, or 3 will have the larger  $K_{eq}$ ? If you do not have enough information to answer this question, write "Can't tell" below. Explain your answer.
- d) Assume that you have a reaction X  $\rightleftharpoons$  Y where  $\Delta G = 0$  and the energy of activation = 0.6 kcal/mole. In a cell, you find that the reaction proceeds almost exclusively in the forward direction. Explain why this might be the case.

MIT OpenCourseWare http://ocw.mit.edu

7.01SC Fundamentals of Biology Fall 2011

For information about citing these materials or our Terms of Use, visit: http://ocw.mit.edu/terms.