In-Class Problems Week 3, Wed.

Problem 1.

Set Formulas and Propositional Formulas.

(a) Verify that the propositional formula $(P \text{ AND } \overline{Q})$ OR (P AND Q) is equivalent to P.

(**b**) Prove that¹

$$A = (A - B) \cup (A \cap B)$$

for all sets, A, B, by showing

 $x \in A$ IFF $x \in (A - B) \cup (A \cap B)$

for all elements, x, using the equivalence of part (a) in a chain of IFF's.

Problem 2.

A *formula of set theory* is a predicate formula that only uses the predicate " $x \in y$." The domain of discourse is the collection of sets, and " $x \in y$ " is interpreted to mean the set x is one of the elements in the set y.

For example, since x and y are the same set iff they have the same members, here's how we can express equality of x and y with a formula of set theory:

$$(x = y) ::= \forall z. (z \in x \text{ IFF } z \in y).$$

$$(1)$$

Express each of the following assertions about sets by a formula of set theory.

(a) $x = \emptyset$.

(b) $x = \{y, z\}.$

(c) $x \subseteq y$. (x is a subset of y that might equal y.)

Now we can explain how to express "x is a proper subset of y" as a set theory formula using things we already know how to express. Namely, letting " $x \neq y$ " abbreviate NOT(x = y), the expression

$$(x \subseteq y \text{ AND } x \neq y),$$

describes a formula of set theory that means $x \subset y$.

From here on, feel free to use any previously expressed property in describing formulas for the following: (d) $x = y \cup z$.

(e) x = y - z.

¹The set difference, A - B, of sets A and B is

$$A - B ::= \{a \in A \mid a \notin B\}.$$

^{2015,} Eric Lehman, F Tom Leighton, Albert R Meyer. This work is available under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 license.

(f) x = pow(y).

(g) $x = \bigcup_{z \in y} z$.

This means that y is supposed to be a collection of sets, and x is the union of all them. A more concise notation for " $\bigcup_{z \in y} z$ ' is simply " $\bigcup y$."

Problem 3.

Forming a pair (a, b) of items a and b is a mathematical operation that we can safely take for granted. But when we're trying to show how all of mathematics can be reduced to set theory, we need a way to represent the pair (a, b) as a set.

(a) Explain why representing (a, b) by $\{a, b\}$ won't work.

(b) Explain why representing (a, b) by $\{a, \{b\}\}$ won't work either. *Hint:* What pair does $\{\{1\}, \{2\}\}$ represent?

(c) Define

$$pair(a, b) ::= \{a, \{a, b\}\}.$$

Explain why representing (a, b) as pair(a, b) uniquely determines a and b. *Hint:* Sets can't be indirect members of themselves: $a \in a$ never holds for any set a, and neither can $a \in b \in a$ hold for any b.

Problem 4.

Subset take-away² is a two player game played with a finite set, A, of numbers. Players alternately choose nonempty subsets of A with the conditions that a player may not choose

- the whole set A, or
- any set containing a set that was named earlier.

The first player who is unable to move loses the game.

For example, if the size of A is one, then there are no legal moves and the second player wins. If A has exactly two elements, then the only legal moves are the two one-element subsets of A. Each is a good reply to the other, and so once again the second player wins.

The first interesting case is when A has three elements. This time, if the first player picks a subset with one element, the second player picks the subset with the other two elements. If the first player picks a subset with two elements, the second player picks the subset whose sole member is the third element. In both cases, these moves lead to a situation that is the same as the start of a game on a set with two elements, and thus leads to a win for the second player.

Verify that when A has four elements, the second player still has a winning strategy.³

²From Christenson & Tilford, David Gale's Subset Takeaway Game, American Mathematical Monthly, Oct. 1997

³David Gale worked out some of the properties of this game and conjectured that the second player wins the game for any set A. This remains an open problem.

6.042J / 18.062J Mathematics for Computer Science Spring 2015

For information about citing these materials or our Terms of Use, visit: https://ocw.mit.edu/terms.