

Thermodynamics of Materials

3.00 Fall 2002

Problem Set 4

Exercise 4.1

Let the molar internal energy and molar entropy of an ideal gas at standard temperature and pressure be $\bar{U}_o(P_o, T_o)$ and $\bar{S}_o(P_o, T_o)$.

Derive expressions for the state functions $\bar{U}(P, T)$, $\bar{U}(V, T)$, $\bar{U}(P, V)$ and $\bar{S}(P, T)$, $\bar{S}(V, T)$, $\bar{S}(P, V)$.

Exercise 4.2

The molar heat capacity of solid aluminum is a weak function of temperature

$$\bar{C}_p = 20.7 + 12.4 \times 10^{-3} T \left(\frac{\text{joule}}{\text{degree mole}} \right) \text{ for } 300\text{K} < T < 900\text{K}.$$

Suppose c grams of aluminum at temperature $T_c = 600(1 - \frac{\theta}{2})$ (kelvin) are put in thermal contact with h grams of aluminum at $T_h = 600(1 + \frac{\theta}{2})$ (kelvin).

- Calculate an expression for the final temperature as a function of the ratio c/h and θ ($0 < \theta < 1$).
- Plot the change in total entropy per mole of aluminum as a function of θ for values of $c/h = (1/16, 1/8, 1/4, 1/2, 1)$.

Hints: Regard the two masses of aluminum together as comprising an isolated system. Work out an algebraic expression for the first part—using a program like Mathematica or Matlab on Athena will make life much much easier. Don't try to calculate and write out an expression for the second part—it is too messy. Find a team member that knows how to plot numbers and make sure that the plot that he or she produces makes sense.