## 18.100B Problem Set 6

Due Friday October 27, 2006 by 3 PM

## **Problems:**

- 1) Prove that if  $\sum |a_n|$  is converges, then  $\sum |a_n|^2$  also converges.
- 2) Prove that

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)(n+2)} = \frac{1}{4}.$$

*Hint:* Use a "telescope trick", i. e. an argument of the form  $\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} (b_n - b_{n+1}) = b_1$ .

- 3) Investigate the behavior (convergence and divergence) of  $\sum_{n=1}^{\infty} a_n$  if
  - a)  $a_n = \sqrt{n+1} \sqrt{n}$ b)  $a_n = \frac{\sqrt{n+1} - \sqrt{n}}{n}$
  - c)  $a_n = \frac{1}{1 + \alpha^n}$  where  $\alpha \ge 0$  is some fixed number.
- 4) Show that convergence of  $\sum_{n=1}^{\infty} a_n$  with  $a_n \ge 0$  implies convergence of

$$\sum_{n=1}^{\infty} \frac{\sqrt{a_n}}{n}.$$

*Hint:* Cauchy–Schwarz inequality.

5) Assume  $a_0 \ge a_1 \ge a_2 \ge \cdots$  and suppose that  $\sum a_n$  converges. Prove that  $\lim_{n \to \infty} (na_n) = 0.$ 

*Hint:* Use and show the inequality  $na_{2n} \leq \sum_{k=n+1}^{2n} a_k$ .

- 6) If X and Y are metric spaces and  $f: X \to Y$  is a mapping between them, show that the following statements are equivalent:
  - a)  $f^{-1}(B)$  is open in X whenever B is open in Y.
  - b)  $f^{-1}(B)$  is closed in X whenever B is closed in Y.
  - c)  $f(\overline{A}) \subseteq \overline{f(A)}$  for every subset A of X.
- 7) Suppose that  $f : \mathbb{R} \to \mathbb{R}$  satisfies

$$\lim_{h \to 0} \left[ f(x+h) - f(x-h) \right] = 0$$

for every  $x \in \mathbb{R}$ . Does this imply that f is continuous?

## Extra problems:

1) Consider the non-absolutetly convergent series

$$\sum_{k=1}^{\infty} a_k = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \frac{1}{6} + \cdots$$

Find a rearrangement  $\sum_{n=1}^{\infty} a_{k_n}$  that diverges to  $+\infty$ .

2) Let  $(p_k) = (2, 3, 5, 7, 11, ...)$  be the sequence of prime numbers. Prove that the series

$$\sum_{k=1}^{\infty} \frac{1}{p_k}$$

diverges, for instance, by working out the details of the following argument.

Suppose  $\sum \frac{1}{p_k}$  converges. Then, there is an integer  $\ell$  such that

$$\sum_{k=\ell+1}^{\infty} \frac{1}{p_k} = \sum_{k=1}^{\infty} \frac{1}{p_{k+\ell}} < \frac{1}{2}$$

Next, for any given  $x \ge 0$ , we let N(x) be the number of positive integers  $n \le x$  that are not divisible by any prime  $p_k$  with  $k > \ell$ . By writing  $n = km^2$  with k square-free (which can be done with any integer), prove that

$$N(x) \le 2^{\ell} \sqrt{x}$$
 and conclude that  $\frac{x}{2} < N(x) < 2^{\ell} \sqrt{x}$ .

This is a contradiction whenever  $x \ge 2^{2\ell+2}$  holds.

Trivia: As an amusing fact, we remark that the series

$$(*) \qquad \sum_{k=1}^{\infty} \left(\frac{1}{p_k} + \frac{1}{q_k}\right) = \left(\frac{1}{3} + \frac{1}{5}\right) + \left(\frac{1}{5} + \frac{1}{7}\right) + \left(\frac{1}{11} + \frac{1}{13}\right) + \left(\frac{1}{17} + \frac{1}{19}\right) + \left(\frac{1}{29} + \frac{1}{31}\right) + \cdots$$

is known to converge. Here  $(p_k, q_k)$  runs over the pairs of *twin primes*; that is,  $p_k$  and  $q_k$  are both prime and  $q_k = p_k + 2$ . This result was proved by V. Brun in 1919, and it is known today (by numerics) that the value of the series is  $\approx 1.902$ .

Had this series (\*) diverged, we would have a proof of the *twin prime conjecture* which means that there are infinitely many twin primes (this is still not known!). Also, if we knew that the value of (\*) were an irrational number, then the twin prime conjecture would follow immediately.

3) Let  $(p_k)$  be the sequence of prime numbers, and let  $J_N$  denote the set of natural numbers whose factorization into primes only involves the primes  $\{p_k : 1 \leq k \leq N\}$ . Prove the following identity

$$\sum_{n \in J_N} \frac{1}{n^s} = \prod_{k=1}^N \frac{1}{1 - p_k^{-s}}$$

for any  $N \in \mathbb{N}$  and any  $s \in \mathbb{Q} \cap (1, \infty)$ . From this deduce Euler's formula

$$\sum_{n=1}^{\infty} \frac{1}{n^s} = \lim_{N \to \infty} \prod_{k=1}^{N} \frac{1}{1 - p_k^{-s}} \Big( =: \prod_{k=1}^{\infty} \frac{1}{1 - p_k^{-s}} \Big)$$

MIT OpenCourseWare http://ocw.mit.edu

18.100B Analysis I Fall 2010

For information about citing these materials or our Terms of Use, visit: http://ocw.mit.edu/terms.