18.440 Final Exam: 100 points
Carefully and clearly show your work on each problem (without writing anything that is technically not true) and put a box around each of your final computations.

- 1. (10 points) Let X be the number on a standard die roll (i.e., each of $\{1,2,3,4,5,6\}$ is equally likely) and Y the number on an independent standard die roll. Write Z=X+Y.
 - 1. Compute the condition probability P[X=4|Z=6].

2. Compute the conditional expectation E[Z|Y] as a function of Y.

- 2. (10 points) Janet is standing outside at time zero when it starts to drizzle. The times at which raindrops hit her are a Poisson point process with parameter $\lambda=2$. In expectation, she is hit by 2 raindrops in each given second.
 - (a) What is the expected amount of time until she is first hit by a raindrop?

(b) What is the probability that she is hit by exactly 4 raindrops during the first 2 seconds of time?

- 3. (10 points) Let X be a random variable with density function f, cumulative distribution function F, variance V and mean M.
 - (a) Compute the mean and variance of 3X + 3 in terms of V and M.

(b) If X_1, \ldots, X_n are independent copies of X. Compute (in terms of F) the cumulative distribution function for the largest of the X_i .

4. (10 points) Suppose that X_i are i.i.d. random variables, each uniform on [0,1]. Compute the moment generating function for the sum $\sum_{i=1}^{n} X_i$.

- 5. (10 points) Suppose that X and Y are outcomes of independent standard die rolls (each equal to $\{1,2,3,4,5,6\}$ with equal probability). Write Z=X+Y.
 - (a) Compute the entropies H(X) and H(Y).

(b) Compute H(X, Z).

(c) Compute H(10X + Y).

(d) Compute $H(Z) + H_Z(Y)$. (Hint: you shouldn't need to do any more calculations.)

- 6. (10 points) Elaine's not-so-trusty old car has three states: broken (in Elaine's possession), working (in Elaine's possession), and in the shop. Denote these states B, W, and S.
 - (i) Each morning the car starts out B, it has a .5 chance of staying B and a .5 chance of switching to S by the next morning.
 - (ii) Each morning the car starts out W, it has .5 chance of staying W, and a .5 chance of switching to B by the next morning.
- (iii) Each morning the car starts out S, it has a .5 chance of staying S and a .5 chance of switching to W by the next morning.

Answer the following

(a) Write the three-by-three Markov transition matrix for this problem.

(b) If the car starts out B on one morning, what is the probability that it will start out B two days later?

(c) Over the long term, what fraction of mornings does the car start out in each of the three states, B, S, and W?

- 7. Suppose that X_1, X_2, X_3, \ldots is an infinite sequence of independent random variables which are each equal to 2 with probability 1/3 and .5 with probability 2/3. Let $Y_0 = 1$ and $Y_n = \prod_{i=1}^n X_i$ for $n \ge 1$.
 - (a) What is the the probability that Y_n reaches 8 before the first time that it reaches $\frac{1}{8}$?

(b) Find the mean and variance of log Y_{10000} .

(c) Use the central limit theorem to approximate the probability that $\log Y_{10000}$ (and hence Y_{10000}) is greater than its median value.

8. (10 points) Eight people toss their hats into a bin and the hats are redistributed, with all of the 8! hat permutations being equally likely. Let N be the number of people who get their own hat. Compute the following:

(a) $\mathbb{E}[N]$

(b) Var[N]

- 9. (10 points) Let X be a normal random variable with mean μ and variance σ^2 .
 - (a) $\mathbb{E}e^X$.

(b) Find μ , assuming that $\sigma^2 = 3$ and $E[e^X] = 1$.

- 10. (10 points)
 - 1. Let $X_1, X_2, ...$ be independent random variables, each equal to 1 with probability 1/2 and -1 with probability 1/2. In which of the cases below is the sequence Y_n a martingale? (Just circle the corresponding letters.)
 - (a) $Y_n = X_n$
 - (b) $Y_n = 1 + X_n$
 - (c) $Y_n = 7$
 - (d) $Y_n = \sum_{i=1}^n iX_i$
 - (e) $Y_n = \prod_{i=1}^n (1 + X_i)$
 - 2. Let $Y_n = \sum_{i=1}^n X_i$. Which of the following is necessarily a stopping time for Y_n ?
 - (a) The smallest n for which $|Y_n| = 5$.
 - (b) The largest n for which $Y_n = 12$ and n < 100.
 - (c) The smallest value n for which n > 100 and $Y_n = 12$.

MIT OpenCourseWare http://ocw.mit.edu

18.440 Probability and Random Variables Spring 2014

For information about citing these materials or our Terms of Use, visit: http://ocw.mit.edu/terms.