18.A34 PROBLEMS #8

- 89. [1] Pat beat Stacy in a set of tennis, winning six games to Stacy's three. Five games were won by the player who did not serve. Who served first?
- 90. [1] Given a seven-minute hourglass and an eleven-minute hourglass, what is the quickest way to time the boiling of an egg for two minutes? Assume that an hourglass cannot be turned upside-down unless it is inactive, i.e., all the sand is at the bottom.
- 91. [1] We all know

$$\int_{-1}^{1} \frac{dx}{1+x^2} = \frac{\pi}{2}.$$

Let x = 1/y, so $dx = -dy/y^2$. When x = 1 and -1, then also y = 1 and -1. Thus

$$\frac{\pi}{2} = \int_{-1}^{1} \frac{-dy/y^2}{1 + (1/y)^2} = -\int_{-1}^{1} \frac{dy}{1 + y^2} = -\frac{\pi}{2}.$$


Hence $\pi = 0$. What (if anything) is wrong?

- 92. [1] Three men play a game with the understanding that the loser is to double the money of the other two. After three games, each has lost just once; and each has \$24. How much did each have to start?
- 93. [1] Move the smallest number of matches to get the cherry outside the glass. The glass may have any orientation at the finish, but of course the cherry cannot be moved.



94. [2] Let n be a positive integer. How many subsets of the set $\{1, 2, ..., n\}$ do not contain two consecutive integers? For instance, when n = 3 there are the five subsets \emptyset , $\{1\}$, $\{2\}$, $\{3\}$, $\{1,3\}$.

- 95. [2] What positive integers can be expressed as the di erence between two squares? The first six are $1 = 1^2 0^2$, $3 = 2^2 1^2$, $4 = 2^2 0^2$, $5 = 3^2 2^2$, $7 = 4^2 3^2$, $8 = 3^2 1^2$.
- 96. [2] In a room 30 feet long, 12 feet wide, and 12 feet high, a fly stands on an end wall one foot above the floor and 6 feet from each side wall. A spider stands on the opposite end wall one foor below the ceiling and 6 feet from each side wall. What is the minimum distance the spider needs to walk along the walls, floor, and ceiling in order to reach the fly?



- 97. (a) [2.5] Can an obtuse triangle be dissected into finitely many acute triangles?
 - (b) [3.5] Can a 3-dimensional cube be dissected into finitely many acute simplices? A (3-dimensional) simplex is any polyhedron with four vertices, six edges, and four triangular faces. A simplex is acute if the angle between any two faces is less than $\pi/2$.
- 98. [2.5] Two baseball teams A and B play against each other for the first time. All that is known about the teams is that A has won a fraction p of its games and that B has won a fraction q, playing against comparable opponents. What is the most reasonable probability f(p,q) that A will beat B? For instance, a little thought will convince you that f(p,1/2) = p, f(p,1) = 0 if $p \neq 1$, f(p,q) + f(q,p) = 1, etc. But it's not so obvious, for instance, what f(.7,.4) should be.
- 99. [2.5] n coins are placed in a circle, either heads or tails up. Player A wants to turn all the coins heads up but is never aware of the status

of the coins. When it is his turn, he can specify any subset of the n positions occupied by the coins. Player B can then rotate the coins around the circle as many spaces as she wants. She then turns over all the coins in the positions specified by A. For what values of n can A win, i.e., in a finite amount of time guarantee that all coins are heads? For instance, A can win when n = 2, by first choosing positions $\{1, 2\}$, then $\{1\}$, and then $\{1, 2\}$. It is easy to check that no matter what the initial configuration of the coins, at some stage both coins will be heads.

100. [3] Let M be the $2^n \times n$ matrix whose rows consist of all 2^n distinct vectors of ± 1 's (in some order) of length n. Change any subset of the entries to 0's. Show that some nonempty subset of the rows of the resulting matrix sums to the vector $[0, 0, \ldots, 0]$.

18.A34 Mathematical Problem Solving (Putnam Seminar) Fall 2018

For information about citing these materials or our Terms of Use, visit: https://ocw.mit.edu/terms.