

MASSACHUSETTS INSTITUTE OF TECHNOLOGY
DEPARTMENT OF PHYSICS
8.962 SPRING 2006

PROBLEM SET 8

Post date: Thursday, April 13th

Due date: Thursday, April 20th

1. In lecture, we derived the following formula for the leading gravitational radiation generated by a source:

$$h_{ij}^{\text{TT}} = \frac{2}{r} \ddot{I}_{kl} \left(P_{ik} P_{jl} - \frac{1}{2} P_{ij} P_{kl} \right) .$$

Here, overdot denotes d/dt , $I_{kl} = \int dV \rho x^k x^l$, we are ignoring the distinction between upstairs and downstairs indices since we are in a nearly flat region and can work in nearly inertial coordinates, and the combination of projection tensors guarantees that the resulting tensor is transverse and traceless.

Show that the same result is obtained if one uses the traceless “quadrupole moment tensor” $\mathcal{I}_{kl} = I_{kl} - \frac{1}{3} \delta_{kl} I$ instead of I_{jk} , where $I = \delta_{kl} I_{kl}$.

Comment 1: This is less trivial than it may seem since there are really two different trace operations defined here: A trace with respect to the spatial metric $\eta_{ij} = \delta_{ij}$, and a trace with respect to the metric of the subspace orthogonal to the propagation direction, P_{ij} .

Comment 2: In general, a radiative l -pole has $2l+1$ separate components — a scalar has one component, a dipole has three, a quadrupole has five. The symmetric spatial tensor I_{ij} has 6 components — there must be “extra” information in that tensor unrelated to radiation. This exercise proves that this extra information is bound up in the trace. For this reason, you will often see the formula for h_{ij}^{TT} written in terms of \mathcal{I}_{ij} rather than I_{ij} .

2. Binary system

Consider a binary consisting of two masses m_1 and m_2 in a circular orbit of radius R about one another. Consider the orbit to be adequately described using Newtonian gravity; in this problem, we will use this description to compute the leading effects due to gravitational-wave emission.

[*Hint:* Don’t forget that orbits in a problem of this type are most easily described using the “reduced system”: a body of mass $\mu = m_1 m_2 / (m_1 + m_2)$ in circular orbit around a body of mass $M = m_1 + m_2$. If you need a refresher, this result is derived in all junior-level mechanics textbooks; see, for example, Goldstein Sec. 3.1.]

(a) Compute the gravitational-wave tensor h_{ij}^{TT} as measured by an observer looking down the angular momentum axis of the system (i.e., the z -axis if you define the orbital plane as the $x - y$ plane).

(b) Compute the rate at which energy is carried away from the system by gravitational waves.

Due to this loss of energy, the radius of the orbit will gradually shrink, and the frequency of the binary will “chirp” to higher frequencies as time passes.

(c) By asserting *global* conservation of energy in the following form,

$$\frac{d}{dt} (E_{\text{kinetic}} + E_{\text{potential}} + E_{\text{GW}}) = 0, \quad (1)$$

derive an equation for dr/dt , the rate at which the orbital radius shrinks.

[*Hint:* Don't forget that for circular, Newtonian orbits, there is a simple relationship expressing $E_{\text{kinetic}} + E_{\text{potential}}$ as a function of r , as well as a simple result for the orbital frequency as a function of r (Kepler's 3rd law).]

(d) Derive the rate of change of the orbital angular frequency Ω due to gravitational-wave emission. You should find that the masses only appear in the combination $\mathcal{M} \equiv \mu^{3/5} M^{2/5}$, perhaps raised to some power. This combination of masses is known as the "chirp mass", since it sets the rate at which the frequency "chirps".

3. Wave equation for the Riemann tensor in linearized theory

As we have emphasized from time to time, there is a nice analogy between the metric of GR and the electromagnetic potential, and likewise between curvature tensors and the electromagnetic field. This suggests that it should be possible to build a wave equation for the curvature tensor.

As background to this problem and the next one, recall that the Einstein field equations can be written in the trace-reversed form

$$R^{\mu}{}_{\alpha\mu\beta} = R_{\alpha\beta} = 8\pi\bar{T}_{\alpha\beta}$$

where $\bar{T}_{\alpha\beta} = T_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}T^{\gamma}{}_{\gamma}$.

To keep things simple, we begin with linearized theory, working in nearly inertial coordinates: $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$ with $||h_{\mu\nu}|| \ll 1$.

(a) To linear order in h , the Bianchi identity can be written

$$\partial_{\alpha}R_{\beta\gamma\mu\nu} + \partial_{\beta}R_{\gamma\alpha\mu\nu} + \partial_{\gamma}R_{\alpha\beta\mu\nu} = 0.$$

Using this equation, show that the divergence of the Riemann tensor is related to the gradient of the trace-reversed stress energy tensor:

$$\partial_{\alpha}R^{\alpha}{}_{\beta\gamma\delta} = 8\pi G \left[\text{source involving gradient of } \bar{T}_{\mu\nu} \right]$$

(b) Now use the Bianchi identity and the solution to part (a) to develop a wave equation for the Riemann tensor of the form

$$\square R_{\alpha\beta\mu\nu} = 8\pi G \left[\text{source involving double gradients of } \bar{T}_{\mu\nu} \right]$$

Solve this equation (formally) using the radiative Green's function introduced in lecture.

(c) Now specialize to a plane gravitational wave propagating in the z -direction through vacuum. The corresponding solution to the above wave equation takes the form $R_{\alpha\beta\mu\nu} = R_{\alpha\beta\mu\nu}(t - z)$. Using the Bianchi identity and the symmetries of Riemann, show that the only non-zero components of the Riemann tensor are of the form R_{i0j0}

(plus components that are trivially related by symmetries; recall that indices i, j only refer to spatial indices).

(d) Show that the only non-zero R_{i0j0} are $R_{x0x0}(t-z) = -R_{y0y0}(t-z)$ and $R_{x0y0}(t-z) = R_{y0x0}(t-z)$. The first non-zero components correspond to the + polarization discussed in lecture; the second corresponds to the \times polarization.

(e) Define fields $h_+(t-z)$ and $h_\times(t-z)$ in terms of these components of the Riemann tensor by

$$R_{x0x0} = -\frac{1}{2}\partial_t^2 h_+, \quad R_{x0y0} = -\frac{1}{2}\partial_t^2 h_\times .$$

Also recall the expression for the Riemann tensor in linearized theory:

$$R_{\alpha\beta\mu\nu} = \frac{1}{2}(\partial_\alpha\partial_\nu h_{\beta\mu} + \partial_\beta\partial_\mu h_{\alpha\nu} - \partial_\alpha\partial_\mu h_{\beta\nu} - \partial_\beta\partial_\nu h_{\alpha\mu})$$

Comparing these two forms, show that $h_+ = h_{xx}^{\text{TT}} = -h_{yy}^{\text{TT}}$, $h_\times = h_{xy}^{\text{TT}} = h_{yx}^{\text{TT}}$.

(f) Show that when one rotates the coordinate system about the waves' propagation direction (i.e., about the z -axis) by an angle θ [so that $x' + iy' = (x + iy)e^{-i\theta}$], the gravitational-wave fields h_+ and h_\times transform such that

$$h'_+ + ih'_\times = (h_+ + ih_\times)e^{-i2\theta}$$

This equation is equivalent to the statement that the graviton is spin 2 — the quantity $h_+ + ih_\times$ has “spin-weight” 2.

4. Nonlinear wave equation for the Riemann tensor

Begin now with the full Bianchi identity:

$$\nabla_\alpha R_{\beta\gamma\mu\nu} + \nabla_\beta R_{\gamma\alpha\mu\nu} + \nabla_\gamma R_{\alpha\beta\mu\nu} = 0 .$$

(a) Develop the fully covariant analog to your answer to part (a) of problem (3):

$$\nabla_\alpha R^\alpha{}_{\beta\gamma\delta} = 8\pi G \left[\text{source involving covariant gradient of } \bar{T}_{\mu\nu} \right]$$

(b) Using the Bianchi identity and the solution to part (a), develop a nonlinear wave equation for the Riemann tensor of the form

$$\square R^\alpha{}_{\beta\gamma\delta} = 8\pi G [\text{source}] .$$

In this case, the source should involve three kinds of terms: double covariant gradients of $\bar{T}_{\mu\nu}$, coupling of $\bar{T}_{\mu\nu}$ to the Riemann tensor, and coupling of Riemann to Riemann. Here, the operator $\square = \nabla^\mu \nabla_\mu$.

Comment: This non-linear wave equation was first developed by Roger Penrose, and is sometimes called the Penrose wave equation. A variant of this equation based on an expansion around the Riemann tensor for a black hole spacetime was developed by Teukolsky and has played an extremely important role in astrophysical relativity research.